


Ordinary Differential Equations

Everything is ordinary about them

<http://nm.mathforcollege.com>

Popping tags means

- A. Popping bubble wrap
- B. Using firecrackers
- C. Changing tags of regular items in a store with tags from clearance items
- D. Taking illicit drugs



<http://nm.mathforcollege.com>

Physical Examples

<http://nm.mathforcollege.com>

How long will it take to cool the trunnion?



$$mc \frac{d\theta}{dt} = -hA(\theta - \theta_a), \quad \theta(0) = \theta_{room}$$

<http://nm.mathforcollege.com>

END

<http://nm.mathforcollege.com>

What did I learn in the ODE class?


<http://nm.mathforcollege.com>

In the differential equation

$$\frac{dy}{dx} + 3y = e^{-x}, y(0) = 6$$

the variable x is the _____ variable

- A. Independent
- B. Dependent




In the differential equation

$$\frac{dy}{dx} + 3y = e^{-x}, y(0) = 6$$

the variable y is the _____ variable

- A. Independent
- B. Dependent




The velocity of a body is given by

$$v(t) = e^{2t} + 5, t \geq 0$$

Then the distance covered by the body from $t=0$ to $t=10$ can be calculated by solving the differential equation for $x(10)$ for


- A. $\frac{dx}{dt} = e^{2t} + 5, x(0) = 0$
- B. $\frac{dx}{dt} = e^{2t} + 5, x(0) = 5$
- C. $\frac{dx}{dt} = 2e^{2t}, x(0) = 0$
- D. $\frac{dx}{dt} = \frac{e^{2t}}{2} + 5t, x(0) = 0$



<http://nm.mathforcollege.com>

The form of the exact solution to $2\frac{dy}{dx} + 3y = e^{-x}, y(0) = 5$ is

- A. $Ae^{-1.5x} + Be^{-x}$
- B. $Ae^{-1.5x} + Bxe^{-x}$
- C. $Ae^{1.5x} + Be^{-x}$
- D. $Ae^{1.5x} + Bxe^{-x}$



<http://nm.mathforcollege.com>

END

<http://nm.mathforcollege.com>

8.03

Euler's Method


<http://nm.mathforcollege.com>

Euler's method of solving ordinary differential equations

$$\frac{dy}{dx} = f(x, y), y(0) = 0 \text{ states}$$

- A. $y_{i+1} = y_i + f(x, y)h$
- B. $y_{i+1} = y_i + f(x_i, y_i)h$
- C. $y_{i+1} = y_i + f(x_i, y_i)$
- D. $y_{i+1} = f(x_i, y_i)h$

<http://nm.mathforcollege.com>




To solve the ordinary differential equation

$$3\frac{dy}{dx} + 5y^2 = \sin x, y(0) = 5,$$

by Euler's method, you need to rewrite the equation as


- A. $\frac{dy}{dx} = \sin x - 5y^2, y(0) = 5$
- B. $\frac{dy}{dx} = \frac{1}{3}(\sin x - 5y^2), y(0) = 5$
- C. $\frac{dy}{dx} = \frac{1}{3}(-\cos x - \frac{5y^3}{3}), y(0) = 5$
- D. $\frac{dy}{dx} = \frac{1}{3}\sin x, y(0) = 5$

<http://nm.mathforcollege.com>



The order of accuracy for a single step in Euler's method is


- A. $O(h)$
- B. $O(h^2)$
- C. $O(h^3)$
- D. $O(h^4)$



<http://nm.mathforcollege.com>

The order of accuracy from initial point to final point while using more than one step in Euler's method is

- A. $O(h)$
- B. $O(h^2)$
- C. $O(h^3)$
- D. $O(h^4)$




<http://nm.mathforcollege.com>

END

<http://nm.mathforcollege.com>

Do you know how Runge- Kutta 4th
Order Method works?

- A. Yes
- B. No
- C. Maybe
- D. I take the 5th



<http://nm.mathforcollege.com>

RUNGE-KUTTA 4TH ORDER METHOD

<http://nm.mathforcollege.com>

Runge-Kutta 4th Order Method

$$\frac{dy}{dx} = f(x, y), y(0) = y_0$$

$$y_{i+1} = y_i + \frac{1}{6}(k_1 + 2k_2 + 2k_3 + k_4)h$$

$$k_1 = f(x_i, y_i)$$

$$k_2 = f\left(x_i + \frac{1}{2}h, y_i + \frac{1}{2}k_1h\right)$$

$$k_3 = f\left(x_i + \frac{1}{2}h, y_i + \frac{1}{2}k_2h\right)$$

$$k_4 = f(x_i + h, y_i + k_3h)$$

<http://nm.mathforcollege.com>

END

<http://nm.mathforcollege.com>

Physical Examples

<http://nm.mathforcollege.com>

Ordinary Differential Equations

Problem:

The trunnion initially at room temperature is put in a bath of dry-ice/alcohol. How long do I need to keep it in the bath to get maximum contraction (“within reason”)?



<http://nm.mathforcollege.com>

Assumptions

The trunnion is a lumped mass system.

- a. What does a lumped system mean? It implies that the internal conduction in the trunnion is large enough that the temperature throughout the ball is uniform.
- b. This allows us to make the assumption that the temperature is only a function of time and not of the location in the trunnion.

<http://nm.mathforcollege.com>

Energy Conservation

$$\text{Heat In} - \text{Heat Lost} = \text{Heat Stored}$$

<http://nm.mathforcollege.com>

Heat Lost

Rate of heat lost
due to convection = $hA(T - T_a)$

h = convection coefficient ($\text{W}/(\text{m}^2 \cdot \text{K})$)

A = surface area, m^2

T = temp of trunnion at a given time, K

<http://nm.mathforcollege.com>

Heat Stored

Heat stored by mass = mCT

where

m = mass of ball, kg

C = specific heat of the ball, J/(kg-K)

<http://nm.mathforcollege.com>

Energy Conservation

Rate at which heat is gained

– Rate at which heat is lost

= Rate at which heat is stored

$$0 - hA(T - T_a) = d/dt(mCT)$$

$$0 - hA(T - T_a) = m C dT/dt$$

<http://nm.mathforcollege.com>

Putting in The Numbers

Length of cylinder = 0.625 m

Radius of cylinder = 0.3 m

Density of cylinder material $\rho = 7800 \text{ kg/m}^3$

Specific heat, $C = 450 \text{ J/(kg-C)}$

Convection coefficient, $h = 90 \text{ W/(m}^2\text{-C)}$

Initial temperature of the trunnion, $T(0) = 27^\circ\text{C}$

Temperature of dry-ice/alcohol, $T_a = -78^\circ\text{C}$

<http://nm.mathforcollege.com>

The Differential Equation

Surface area of the trunnion

$$\begin{aligned} A &= 2\pi rL + 2\pi r^2 \\ &= 2 * \pi * 0.3 * 0.625 + 2 * \pi * 0.3^2 \\ &= 1.744 \text{ m}^2 \end{aligned}$$

Mass of the trunnion

$$\begin{aligned} M &= \rho V \\ &= \rho (\pi r^2 L) \\ &= (7800) * [\pi * (0.3)^2 * 0.625] \\ &= 1378 \text{ kg} \end{aligned}$$

<http://nm.mathforcollege.com>

The Differential Equation

$$-hA(T - T_a) = mC \frac{dT}{dt}$$


$$-90 \times 1.744 \times (T + 78) = 1378 \times 450 \frac{dT}{dt}$$

$$\frac{dT}{dt} = -2.531 \times 10^{-4} (T + 78),$$

$$T(0) = 27$$

<http://nm.mathforcollege.com>

Solution



Time (s)	Temp (°C)
0	27
1000	0.42
2000	-19.42
3000	-34.25
4000	-45.32
5000	-53.59
6000	-59.77
7000	-64.38
8000	-67.83
9000	-70.40
10000	-72.32

<http://nm.mathforcollege.com>


END

<http://nm.mathforcollege.com>

Classify the differential equation

$$2 \frac{dy}{dx} + 4y = e^x + 3, y(0) = 5$$

- A. linear
- B. nonlinear
- C. undeterminable to be linear or nonlinear




<http://nm.mathforcollege.com>

Classify the differential equation

$$2 \frac{dy}{dx} + xy = e^x + 3, y(0) = 5$$

- A. linear
- B. nonlinear
- C. linear with fixed constants
- D. undeterminable to be linear or nonlinear




<http://nm.mathforcollege.com>

Classify the differential equation

$$2 \frac{dy}{dx} + y^2 = e^x + 3, y(0) = 5$$


- A. linear
- B. nonlinear
- C. linear with fixed constants
- D. undeterminable to be linear or nonlinear



<http://nm.mathforcollege.com>

Ordinary differential equations can have these many **dependent** variables.

- A. one
- B. two
- C. any positive integer



Ordinary differential equations can have these many **independent** variables.

- A. one
- B. two
- C. any positive integer

