Special Chapter 04.09 Adequacy of Solutions

After reading this chapter, you should be able to:

- 1. differentiate between ill-conditioned and well-conditioned systems of equations,
- 2. define and find the norm of a matrix,
- 3. define and evaluate the condition number of an invertible square matrix,
- 4. relate the condition number of a coefficient matrix to the ill or well conditioning of the system of simultaneous linear equations, that is, how much can you trust the solution of the simultaneous linear equations.

What do you mean by an ill-conditioned and a well-conditioned system of equations?

A system of equations is considered to be **well-conditioned** if a small change in the coefficient matrix or a small change in the right-hand side results in a small change in the solution vector.

A system of equations is considered to be **ill-conditioned** if a small change in the coefficient matrix or a small change in the right-hand side results in a large change in the solution vector.

Example 1

Is this system of equations well-conditioned?

$$\begin{bmatrix} 1 & 2 \\ 2 & 3.999 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 7.999 \end{bmatrix}$$

Solution

The solution to the above set of equations is

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

Make a small change in the right-hand side vector of the equations

$$\begin{bmatrix} 1 & 2 \\ 2 & 3.999 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4.001 \\ 7.998 \end{bmatrix}$$

gives

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -3.999 \\ 4.000 \end{bmatrix}$$

Make a small change in the coefficient matrix of the equations

$$\begin{bmatrix} 1.001 & 2.001 \\ 2.001 & 3.998 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 7.999 \end{bmatrix}$$

gives

Page 1 of 12

 $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 3.994 \\ 0.001388 \end{bmatrix}$

This last system of equations "looks" ill-conditioned because a small change in the coefficient matrix or the right-hand side resulted in a large change in the solution vector.

Example 2

Is this system of equations well-conditioned?

 $\begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 7 \end{bmatrix}$

Solution

The solution to the above equations is

 $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$

Make a small change in the right-hand side vector of the equations.

 $\begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4.001 \\ 7.001 \end{bmatrix}$

gives

 $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1.999 \\ 1.001 \end{bmatrix}$

Make a small change in the coefficient matrix of the equations.

1.001]	ן2.001	$[x]_{-}$	_ [4]
l2.001	3.001 []]	$\lfloor y \rfloor^-$	- [2]

gives

 $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2.003 \\ 0.997 \end{bmatrix}$

This system of equations "looks" well-conditioned because small changes in the coefficient matrix or the right-hand side resulted in small changes in the solution vector.

So, what if the system of equations is ill-conditioned or well-conditioned?

Well, if a system of equations is ill-conditioned, we cannot trust the solution as much. Revisit the velocity problem, Example 1 in Chapter 04.06. The values in the coefficient matrix [A] are squares of time, etc. For example, if instead of $a_{11} = 25$, you used $a_{11} = 24.99$, would you want this small change to make a huge difference in the solution vector. If it did, would you trust the solution?

04.09.2

Later we will see how much (quantifiable terms) we can trust the solution in a system of equations. Every invertible square matrix has a **condition number** and coupled with the **machine epsilon**, we can quantify how many significant digits one can trust in the solution.

To calculate the condition number of an invertible square matrix, I need to know what the norm of a matrix means. How is the norm of a matrix defined?

Just like the determinant, the norm of a matrix is a simple unique scalar number. However, the norm is always positive and is defined for all matrices – square or rectangular, and invertible or noninvertible square matrices.

One of the popular definitions of a norm is the row sum norm (also called the uniform-matrix norm). For a $m \times n$ matrix [A], the row sum norm of [A] is defined as

$$\|A\|_{\infty} = \max_{1 \le i} \sum_{j=1}^{n} |a_{ij}|$$

that is, find the sum of the absolute value of the elements of each row of the matrix [A]. The maximum out of the *m* such values is the row sum norm of the matrix [A].

Example 3

Find the row sum norm of the following matrix [A].

$$A = \begin{bmatrix} 10 & -7 & 0 \\ -3 & 2.099 & 6 \\ 5 & -1 & 5 \end{bmatrix}$$

Solution

$$\|A\|_{\infty} = \max_{1 \le i \le 3} \sum_{j=1}^{3} |a_{ij}|$$

= max[(|10| + |-7| + |0|), (|-3| + |2.099| + |6|), (|5| + |-1| + |5|)]
= max[(10 + 7 + 0), (3 + 2.099 + 6), (5 + 1 + 5)]
= max[17, 11.099, 11]
= 17

How is the norm related to the conditioning of the matrix?

Let us start answering this question using an example. Go back to the *ill-conditioned* system of equations,

 $\begin{bmatrix} 1 & 2 \\ 2 & 3.999 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 7.999 \end{bmatrix}$

that gives the solution as

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

Denoting the above set of equations as

[A][X] = [C] $||X||_{\infty} = 2$ $||C||_{\infty} = 7.999$

Making a small change in the right-hand side,

$$\begin{bmatrix} 1 & 2 \\ 2 & 3.999 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4.001 \\ 7.998 \end{bmatrix}$$

gives

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} -3.999 \\ 4.000 \end{bmatrix}$$

Denoting the above set of equations by

[A][X'] = [C']

right-hand side vector is found by

 $[\Delta C] = [C'] - [C]$

and the change in the solution vector is found by

$$[\Delta X] = [X'] - [X]$$

then

$$\begin{bmatrix} \Delta C \end{bmatrix} = \begin{bmatrix} 4.001 \\ 7.998 \end{bmatrix} - \begin{bmatrix} 4 \\ 7.999 \end{bmatrix}$$
$$= \begin{bmatrix} 0.001 \\ -0.001 \end{bmatrix}$$

and

$$\begin{bmatrix} \Delta X \end{bmatrix} = \begin{bmatrix} -3.999\\ 4.000 \end{bmatrix} - \begin{bmatrix} 2\\ 1 \end{bmatrix}$$
$$= \begin{bmatrix} -5.999\\ 3.000 \end{bmatrix}$$

then

 $\|\Delta C\|_{\infty}=0.001$

 $\|\Delta X\|_{\infty} = 5.999$

The relative change in the norm of the solution vector is

$$\frac{\|\Delta X\|_{\infty}}{\|X\|_{\infty}} = \frac{5.999}{2} = 2.9995$$

The relative change in the norm of the right-hand side vector is

$$\frac{\|\Delta C\|_{\infty}}{\|C\|_{\infty}} = \frac{0.001}{7.999}$$
$$= 1.250 \times 10^{-4}$$

See the small relative change of 1.250×10^{-4} in the right-hand side vector norm results in a large relative change in the solution vector norm of 2.9995.

In fact, the ratio between the relative change in the norm of the solution vector and the relative change in the norm of the right-hand side vector is

$$\frac{\|\Delta X\|_{\infty}/\|X\|_{\infty}}{\|\Delta C\|_{\infty}/\|C\|_{\infty}} = \frac{2.9995}{1.250 \times 10^{-4}} = 23993$$

Let us now go back to the *well-conditioned* system of equations.

$$\begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 7 \end{bmatrix}$$

gives

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 \\ 1 \end{bmatrix}$$

Denoting the system of equations by

$$[A][X] = [C]$$
$$\|X\|_{\infty} = 2$$
$$\|C\|_{\infty} = 7$$

Making a small change in the right-hand side vector

$$\begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4.001 \\ 7.001 \end{bmatrix}$$

gives

 $\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 1.999 \\ 1.001 \end{bmatrix}$ Denoting the above set of equations by

$$[A][X'] = [C']$$

the change in the right-hand side vector is then found by

$$[\Delta C] = [C'] - [C]$$

and the change in the solution vector is

$$[\Delta X] = [X'] - [X]$$

then

$$\begin{bmatrix} \Delta C \end{bmatrix} = \begin{bmatrix} 4.001 \\ 7.001 \end{bmatrix} - \begin{bmatrix} 4 \\ 7 \end{bmatrix}$$
$$= \begin{bmatrix} 0.001 \\ 0.001 \end{bmatrix}$$

and

$$\begin{bmatrix} \Delta X \end{bmatrix} = \begin{bmatrix} 1.999\\ 1.001 \end{bmatrix} - \begin{bmatrix} 2\\ 1 \end{bmatrix}$$
$$= \begin{bmatrix} -0.001\\ 0.001 \end{bmatrix}$$

then

$$\|\Delta C\|_{\infty} = 0.001$$
$$\|\Delta X\|_{\infty} = 0.001$$

The relative change in the norm of solution vector is

$$\frac{\|\Delta X\|_{\infty}}{\|X\|_{\infty}} = \frac{0.001}{2} = 5 \times 10^{-4}$$

The relative change in the norm of the right-hand side vector is

$$\frac{\|\Delta C\|_{\infty}}{\|C\|_{\infty}} = \frac{0.001}{7}$$
$$= 1.429 \times 10^{-4}$$

See the small relative change in the right-hand side vector norm of 1.429×10^{-4} results in the small relative change in the solution vector norm of 5×10^{-4} .

In fact, the ratio between the relative change in the norm of the solution vector and the relative change in the norm of the right-hand side vector is

$$\frac{\|\Delta X\|_{\infty}/\|X\|_{\infty}}{\|\Delta C\|_{\infty}/\|C\|_{\infty}} = \frac{5 \times 10^{-4}}{1.429 \times 10^{-4}} = 3.5$$

What are some of the properties of norms?

- 1. For a matrix [A], $||A|| \ge 0$
- 2. For a matrix [A] and a scalar k, ||kA|| = |k|||A||
- 3. For two matrices [A] and [B] of the same order, $||A + B|| \le ||A|| + ||B||$
- 4. For two matrices [A] and [B] that can be multiplied as [A] [B], $||AB|| \le ||A|| ||B||$

Is there a general relationship that exists between $||\Delta X||/||X||$ and $||\Delta C||/||C||$ or between $||\Delta X||/||X||$ and $||\Delta A||/||A||$? If so, it could help us identify a well-conditioned and an ill-conditioned system of equations.

If there is such a relationship, will it help us quantify the conditioning of the matrix? That is, will it tell us how many significant digits we could trust in the solution of a system of simultaneous linear equations?

There is a relationship that exists between

$$\frac{\|\Delta X\|}{\|X\|} \text{ and } \frac{\|\Delta C\|}{\|C\|}$$

and between

$$\frac{\|\Delta X\|}{\|X\|} \text{ and } \frac{\|\Delta A\|}{\|A\|}$$

These relationships are

$$\frac{\|\Delta X\|}{\|X\|} \le \|A\| \|A^{-1}\| \frac{\|\Delta C\|}{\|C\|}$$

and

$$\frac{\|\Delta X\|}{\|X + \Delta X\|} \le \|A\| \|A^{-1}\| \frac{\|\Delta A\|}{\|A\|}$$

The above two inequalities show that the relative change in the norm of the right-hand side vector or the coefficient matrix can be amplified by as much as $||A|| ||A^{-1}||$.

This number $||A|| ||A^{-1}||$ is called the **condition number** of the matrix and coupled with the machine epsilon, we can quantify the accuracy of the solution of [A] [X] = [C].

How do I use the above theorems to find how many significant digits are correct in my solution vector?

The relative error in a solution vector norm is \leq Cond(A) × relative error in the right-hand side vector norm.

The possible relative error in the solution vector norm is $\leq Cond(A) \times \epsilon_{mach}$

Hence Cond(A) $\times \in_{mach}$ should give us the number of significant digits, *m* that are at least correct in our solution by finding out the largest value of *m* for which Cond(A) $\times \in_{mach}$ is less than 0.5×10^{-m} .

Example 4

How many significant digits can I trust in the solution of the following system of equations? Assume single precision for real numbers where 32 bits are used to represent a number, with 1 bit for the sign of the number, 7 bits for the biased exponent, and 23 bits for the magnitude of the mantissa.

$$\begin{bmatrix} 1 & 2 \\ 2 & 3.999 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 2 \\ 4 \end{bmatrix}$$

Solution

For

$$[A] = \begin{bmatrix} 1 & 2 \\ 2 & 3.999 \end{bmatrix}$$

it can be shown

$$[A]^{-1} = \begin{bmatrix} -3999 & 2000 \\ 2000 & -1000 \end{bmatrix}$$
$$\|A\|_{\infty} = 5.999$$
$$\|A^{-1}\|_{\infty} = 5999$$
$$Cond(A) = \|A\| \|A^{-1}\|$$
$$= 5.999 \times 5999.4$$
$$= 35990$$

The machine epsilon is

$$\epsilon_{mach} = 2^{-23}$$

= 0.119209 × 10⁻⁶

 $Cond(A) \times \in_{mach} = 35990 \times 0.119209 \times 10^{-6}$ = 0.4290 × 10⁻²

For what maximum positive value of *m*, would $Cond(A) \times \in_{mach}$ be less than or equal to 0.5×10^{-m}

 $\begin{array}{l} 0.4290 \times 10^{-2} \leq 0.5 \times 10^{-m} \\ 0.8580 \times 10^{-2} \leq 10^{-m} \\ \log(0.8580 \times 10^{-2}) \leq \log(10^{-m}) \\ -2.067 \leq -m \\ m \leq 2.067 \\ m \leq 2 \end{array}$

So, two significant digits are at least correct in the solution vector.

Example 5

How many significant digits can I trust in the solution of the following system of equations?

$$\begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} 4 \\ 7 \end{bmatrix}$$

Solution

For

$$[A] = \begin{bmatrix} 1 & 2 \\ 2 & 3 \end{bmatrix}$$

it can be shown

$$[A]^{-1} = \begin{bmatrix} -3 & 2\\ 2 & -1 \end{bmatrix}$$

Then

$$||A||_{\infty} = 5,$$

$$||A^{-1}||_{\infty} = 5.$$

Cond (A) = ||A|| ||A^{-1}||
= 5 × 5
= 25

Assuming single precision with 23 bits used in the mantissa for real numbers, the machine epsilon

 $\in_{mach} = 2^{-23}$

04.09.10

 $= 0.119209 \times 10^{-6}$ Cond(A) × ϵ_{mach} = 25 × 0.119209 × 10⁻⁶ = 0.2980 × 10⁻⁵

For what maximum positive value of *m*, would $Cond(A) \times \in_{mach}$ be less than or equal to 0.5×10^{-m}

 $0.2980 \times 10^{-5} \le 0.5 \times 10^{-m}$

 $m \leq 5$

So, five significant digits are at least correct in the solution vector.

Key Terms:

Ill-Conditioned matrix Well-Conditioned matrix Norm Condition Number Machine Epsilon Significant Digits

Problem Set

1. The row sum norm of the matrix

$$[A] = \begin{bmatrix} 6 & -7 & 3 & 13\\ 19 & -21 & 23 & -29\\ 41 & 47 & -51 & 61 \end{bmatrix}$$

- is (A) 29 (B) 61
- (C) 98
- (D) 200
- 2. The adequacy of the solution of simultaneous linear equations [A][X] = [C] depends on (A) the condition number of coefficient matrix [A]
 - (B) the machine epsilon
 - (C) the condition number for matrix [A] and the machine epsilon
 - (D) norm of the coefficient matrix [A]
- 3. Given a set of equations in matrix form [A][X] = [C], ||A|| = 250, $||A^{-1}|| = 40$ and $\varepsilon_{mach} = 0.119 \times 10^{-6}$, then the number of significant digits you can at least trust in the solution are
 - (A) 1
 - (B) 2
 - (C) 3
 - (D)4
- 4. The condition number of the $n \times n$ identity matrix based on the row sum norm is (A)0
 - (B) 1
 - $(\mathbf{D})\mathbf{1}$ $(\mathbf{C})\mathbf{n}$
 - (C) n^{2}
 - (D)*n*
- 5. For

$$[A] = \begin{bmatrix} 10 & -7 & 0 \\ -3 & 2.099 & 6 \\ 5 & -1 & 5 \end{bmatrix}$$

gives

$$[A]^{-1} = \begin{bmatrix} -0.1099 & -0.2333 & 0.2799 \\ -0.2999 & -0.3332 & 0.3999 \\ 0.04995 & 0.1666 & 6.664 \times 10^{-5} \end{bmatrix}$$

- (A) What is the condition number of [A]?
- (B) How many significant digits can we at least trust in the solution of [A] [X] = [C] if $\epsilon_{mach} = 0.1192 \times 10^{-6}$?